MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2019

April–June 2019—Volume 134, pp. 369–376

https://doi.org/10.5248/134.369

Heteroplacidium compactum reported as a genus new to China

Xiang-Min Cheng[1,2], Da-Le Liu[1,3], Xin-Li Wei*[1,2], Jiang-Chun Wei[1,2]

[1] *State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China*

[2] *University of Chinese Academy of Sciences, Beijing 100049, China*

[3] *College of Life Sciences, Southwest Forestry University, Kunming 650224, China*

* Correspondence to: *weixl@im.ac.cn*

Abstract—Three specimens of *Heteroplacidium compactum* (*Verrucariaceae*), isolated from arid areas in Northwest China, were identified by morphological and phylogenetic comparisons with other species, including species of *Endocarpon* and *Placidium* (common verrucariaceous genera in the same area). This is the first report of *Heteroplacidium* from China.

Key words—desert, ITS, lichen, morphology, taxonomy

Introduction

Breuss (1996) separated four genera—*Anthracocarpon*, *Clavascidium*, *Heteroplacidium*, and *Involucropyrenium*—from *Catapyrenium* Flot. s. lat. (*Verrucariaceae*) based on morphological characters. *Neocatapyrenium* H. Harada, *Placidium* A. Massal., and *Scleropyrenium* H. Harada are also morphologically related (Prieto & al. 2010b) but are independent monophyletic lineages within *Verrucariaceae*, according to phylogenetic analyses (Gueidan & al. 2007, 2009; Prieto & al. 2010b).

Endocarpon Hedw. (*Verrucariaceae*) was relatively common during our field survey in the arid region of Northwest China (Yang & Wei 2008, Zhang & al. 2017a), but no species of *Heteroplacidium* Breuss was recorded. *Heteroplacidium,* which grows primarily on soil and rock in warm temperate



Fig. 1. *Heteroplacidium compactum* collection sites in China.

regions, is characterized by a crustose-areolate to squamulose thallus, a paraplectenchymatous or subparaplectenchymatous cortex, and clavate asci with biseriate ascospores (Prieto & al. 2010a); twelve species have been accepted (Knudsen & al. 2014). We report *Heteroplacidium* in China for the first time, based on three specimens of *H. compactum*.

Material & methods

Phenotypic analysis

All *Heteroplacidium* samples were collected from Ningxia Hui Autonomous Region and Inner Mongolia (Fig. 1) and are preserved in the Lichen Section of Herbarium Mycologicum Academiae Sinicae, Beijing, China (HMAS-L). Zeiss Stemi SV11 dissecting and Zeiss Axioskop 2 plus compound microscopes were used for morphological examinations.

DNA extraction, amplification, sequencing

DNA was extracted from our three fresh specimens of *Heteroplacidium* (Table 1) following the modified CTAB method (Rogers & Bendich 1988). The internal transcribed spacer of nuclear ribosome DNA (nrDNA ITS), the standard DNA barcoding marker (Schoch & al. 2012), was chosen as the genetic marker. Primers LR1 (Vilgalys & Hester 1990) and ITS1 (White & al. 1990) were used. Reactions were carried out in 50 µL reaction volume comprising 3 µL total DNA, 1 µL each primer (10 µM), 25 µL 2×Taq MasterMix (CWBIO) and 20 µL ddH_2O. The DNA was amplified in a Biometra T-Gradient thermal cycler as follows: initial heating for 5 min at 95 °C; 35 cycles of 30 s at 94 °C, 30 s at 56 °C, and 90 s at 72 °C; ending with a final extension of 8 min at 72 °C, after which the samples were kept at 4 °C. Negative controls were prepared for each amplification series. PCR products were purified using a CWBIO gel extraction kit following the manufacturer's instructions.

Sequence alignment and phylogenetic tree construction

PCR products from the three putative *Heteroplacidium* specimens were sequenced using ABI 3730 XL Sequencer. Additional sequences of *Endocarpon* (3 spp.), *Heteroplacidium* (4 spp.), and *Placidium* (3 spp.) were downloaded from GenBank (Table 1) with the *Endocarpon* sequences selected as outgroup. The sequences were aligned using ClustalW Multiple Alignment (Thompson & al. 1994) in BioEdit 7.2.5 (Hall 1999). The program Gblocks v0.91b (Castresana 2000, Talavera & Castresana 2007) was used to delimit and remove regions of alignment uncertainty, using options for a "less stringent" selection on the Gblocks web server (http://molevol.cmima.csic.es/castresana/Gblocks_server.html). We confirmed the phylogenetic position of *Heteroplacidium* sp. represented by our three specimens by subjecting the alignment to a maximum likelihood (RAxML) analysis; nodal support was assessed using 1000 bootstrapping pseudoreplicates with RAxML-HPC v. 8.2.6 (Stamatakis 2014) and MrBayes v.3.2.6 (Huelsenbeck & Ronquist 2001, Ronquist & Huelsenbeck 2003) on the CIPRES Science Gateway (http://www.phylo.org). In the ML and Bayesian analyses, substitution models for ITS were estimated using jModelTest-2.1.9 (Guindon & Gascuel 2003, Darriba & al. 2012). Based

Table 1. Specimens and sequences of *Endocarpon*, *Heteroplacidium*, and *Placidium* used in the phylogenetic analysis

Species	Voucher	Origin	GenBank No.
E. adscendens	CG671	Switzerland	KF959777
E. pusillum	CG470	—	JQ927447
E. tenuissimum	Lendemer 29447	USA	KM371592
	Lendemer 27013	USA	KM371593
H. acervatum	LI 271015	Spain	GU228954
	M. Prieto 399	Spain	GU228955
H. compactum	M. Prieto 1701	Spain	GU228952
	M. Prieto 1607	Spain	GU228949
	XL2017117	China	MH930459*
	ALS2018002	China	MH930460*
	ALS2018036	China	MH930461*
H. congestum	LI 552268	USA	GU228950
	LI 297536	USA	GU228951
H. divisum	LI 218428	Italy	GU228953
P. pilosellum	M. Prieto 3	Spain	GU228993
	VEL_PLG	Slovakia	KY981585
P. podolepis	LI 297365	Argentina	GU228956
P. squamulosum	M. Prieto 336	Spain	GU228994

* = sequences newly generated for this study by the authors

on these results, we used the TrN+I+G model with 1000 pseudoreplicates in the ML analysis, and two parallel Markov chain Monte Carlo (MCMC) runs were performed in MrBayes, each using 8 million generations and sampling every 1000 steps. A 50% majority-rule consensus tree was generated from the combined sampled trees of both runs after discarding the first 25% as burn-in. Tree files were visualized with FigTree v.1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/).



Fig. 2. The maximum likelihood tree of *Endocarpon*, *Heteroplacidium*, and *Placidium* species based on ITS sequences. The numbers in each node represent bootstrap support (BS) and posterior probability (PP) values. Bootstrap values ≥75 and posterior probability values ≥0.95 were plotted on the branches of the RAxML tree. Except for the three samples of *Heteroplacidium compactum* marked by the solid circle '●', all other sequences were downloaded from GenBank. Scale bar = 0.07 substitution per site.

Phylogenetic and species delimitation analyses

The aligned matrix contained 455 unambiguous ITS nucleotide positions. The phylogenetic tree (Fig. 2) included 10 taxa representing three genera within *Verrucariaceae*. *Heteroplacidium* formed a well-supported monophyletic clade (BS = 94, PP = 0.99) with the distinct species clearly separated. Our three Chinese specimens all grouped with Spanish sequences of *H. compactum*.

Taxonomy

Heteroplacidium compactum (A. Massal.) Gueidan & Cl. Roux,
Bull. Inf. Ass. Franç. Lichén. 33(1): 25, 2008. Fig. 3

Thallus areolate, areoles 0.6–1.8 mm diam., angular to rounded, flat to slightly convex; upper surface dark brown to black, dull to somewhat shiny.

Thallus layer ≤300 µm thick; upper cortex paraplectenchymous, 12.5–37.5 µm thick; algal layer 62.5–150 µm thick, cells 12.5–17.5 µm diam; medulla 112.5–187.5 µm thick; lower cortex paraplectenchymous, 10–12.5 µm thick or not clearly delimited. Rhizohyphae simple, intersecting,



Fig. 3. *Heteroplacidium compactum* (HMAS-L 140900): A. Habit of thallus; B. Anatomical structure of thallus; C. Perithecia in thallus; D. Ascospores; E. Separate areole with abundant rhizines tangling with soil particles. Scale bars: A = 1 mm; B = 20 µm; C = 100 µm; D = 50 µm; E = 0.5 mm.

4.5–7.5 µm diam. Perithecia nearly spherical, 300–500 µm diam, exciple, pale darkening with age. Asci clavate, 52.5–62.5 × 15–17.5 µm. Ascospores simple, ellipsoid, 12.5–17.5 × 10–12.5 µm. Pycnidia not seen.

Specimens examined. **CHINA**. **Inner Mongolia**: 39°28′22″N 101°04′04″E, 1563 m alt., on sand, 5 June 2018, D.L. Liu & al. ALS2018002 (HMAS-L 0141672; GenBank MH930460); 39°32′31″N 101°06′34″E, 1478 m alt., on sand, 5 June 2018, D.L. Liu et al. ALS2018036 (HMAS-L 0141671; GenBank MH930461). **Ningxia**: **Zhongwei City**, Cui Liu Gou. 37°24′35″N 104°35′09″E, 1577 m alt., on sand, 15 July 2017, D.L. Liu & R.D. Liu XL2017117 (HMAS-L 140900; GenBank MH930459).

Distribution: *Heteroplacidium compactum* has previously been reported from Asia (Mongolian People's Republic; Pakistan), Europe, and northern Africa (Breuss 1994; Prieto & al. 2010a). In China, it is now known from arid desert areas in Inner Mongolia and Ningxia.

Discussion

The characters of Chinese *H. compactum* are essentially similar to the previous descriptions (Breuss 1994; Prieto & al. 2010a) in the areolate thallus, dark brown upper surface, paraplectenchymous upper and lower cortex, and the shape and size of the perithecia and asci. Among the differences noted are a thinner thallus (300 µm versus 600 µm) and broader ascospores (10–12.5 µm versus 8-10 µm). Furthermore, pycnidia were not seen in the Chinese materials.

In the arid desert area of China, *Endocarpon* is quite common and comprises a key component of biological soil crust communities (BSC). The biodiversity reported for these Chinese BSCs includes several newly described *Endocarpon* species (Yang & Wei 2008, Zhang & al. 2017a) with the predominant species, *E. pusillum*, studied physiologically (Zhang & Wei 2011), ecologically (Ding & al. 2013), and genomically (Wang & al. 2014, 2015, Li & Wei 2016, Zhang & al. 2017b). *Heteroplacidium* (with *Dermatocarpon*-type pycnidia) is clearly different from *Endocarpon* (with *Endocarpon*-type pycnidia).

The most similar genus to *Heteroplacidium* is *Placidium* A. Massal., which is also characterized by *Dermatocarpon*-type pycnidia, but the two genera have differently sized thallus squamules (*Heteroplacidium* 0.3–3 mm diam; *Placidium* 2–8(–15) mm diam). In the phylogenetic tree (Fig. 2), *Heteroplacidium* is most closely related to *Placidium*, clustering in a well-supported clade (BS = 100, PP = 1) and relatively distant from *Endocarpon*.

Acknowledgments

This research was funded by one of National Key Research and Development Program of China (2016YFE0203400) and the National Natural Science Foundation of China (31770022, 31470149). The authors sincerely thank the two peer reviewers, Dr. Steve Leavitt (Brigham Young University, USA) and Dr. Ze-Feng Jia (Liaocheng University, China), for giving constructive suggestions and kindly help in making careful modifications. Ms. H. Deng gave considerable assistance during our studies in HMAS-L.

Literature cited

Breuss O. 1994. Über einige wenig bekannte *Verrucaria*-Arten (Lichenes, *Verrucariaceae*). Österreichische Zeitschrift für Pilzkunde 3: 15–20.

Breuss O. 1996. Ein verfeinertes Gliederungskonzept für *Catapyrenium* (lichenisierte Ascomyceten, *Verrucariaceae*). Annalen Naturhistorischen Museums in Wien 98 B Suppl.: 35–50.

Castresana J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Molecular Biology and Evolution 17(4): 540–552. https://doi.org/10.1093/oxfordjournals.molbev.a026334

Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. Nature Methods 9(8): 772. https://doi.org/10.1038/nmeth.2109

Ding LP, Zhou QM, Wei JC. 2013. Estimation of *Endocarpon pusillum* carbon budget in the Tengger Desert based on its photosynthetic rate. Science China Life Sciences 56(9): 848–855. https://doi.org/10.1007/s11427-013-4526-9

Gueidan C, Roux C, Lutzoni F. 2007. Using a multigene phylogenetic analysis to assess generic delineation and character evolution in *Verrucariaceae* (*Verrucariales*, *Ascomycota*). Mycological Research 111(10): 1145–1168. https://doi.org/10.1016/j.mycres.2007.08.010

Gueidan C, Savić S, Thüs H, Roux C, Keller C, Tibell L, Prieto M, Heiðmarsson S, Breuss O, Orange A, Fröberg L, Wynns AA, Navarro-Rosinés P, Krzewicka B, Pykälä J, Grube M, Lutzoni F. 2009. Generic classification of the *Verrucariaceae* (*Ascomycota*) based on molecular and morphological evidence: recent progress and remaining challenges. Taxon 58: 184–208. https://doi.org/10.1002/tax.581019

Guindon S, Gascuel O. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic Biology 52(5): 696–704. https://doi.org/10.1080/10635150390235520

Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Serie 41: 95–98.

Huelsenbeck JP, Ronquist F. 2001. MrBayes: Bayesian inference of phylogenetic trees. Bioinformatics 17(8): 754–755. https://doi.org/10.1093/bioinformatics/17.8.754

Knudsen K, Breuss O, Kocourková J. 2014. A new lichenicolous *Heteroplacidium* (*Verrucariaceae*) from the deserts of southern California. Opuscula Philolichenum 13: 26–33.

Li H, Wei JC. 2016. Functional analysis of thioredoxin from the desert lichen-forming fungus, *Endocarpon pusillum* Hedwig, reveals its role in stress tolerance. Scientific Reports 6: 27184. https://doi.org/10.1038/srep27184

Prieto M, Aragón G, Martínez I. 2010a. The genus *Catapyrenium* s. lat. (*Verrucariaceae*) in the Iberian Peninsula and the Balearic Islands. Lichenologist 42(6): 637–684. https://doi.org/10.1017/S0024282910000319

Prieto M, Martínez I, Aragón G, Otálora MAG, Lutzoni F. 2010b. Phylogenetic study of *Catapyrenium* s. str. (*Verrucariaceae,* lichen-forming *Ascomycota*) and related genus *Placidiopsis*. Mycologia 102(2): 291–304. https://doi.org/10.3852/09-168

Rogers SO, Bendich AJ. 1988. Extraction of DNA from plant tissues. A6: 1–10, in: Gelvin SB, Schilperoort RA (eds). Plant Molecular Biology Manual. Kluwer Academic Publishers, Boston. https://doi.org/10.1007/978-94-017-5294-7_6

Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574. https://doi.org/10.1093/bioinformatics/btg180

Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA & al. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. Proceedings of the National Academy of Sciences of the United States of America 109(16): 6241–6246. https://doi.org/10.1073/pnas.1117018109

Stamatakis A. 2014. RAxML Version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Talavera G, Castresana J. 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. Systematic Biology 56(4): 564–577. https://doi.org/10.1080/10635150701472164

Thompson J, Higgins D, Gibson T. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Research 22(22): 4673–4680. https://doi.org/10.1093/nar/22.22.4673

Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172: 4238–4246. https://doi.org/10.1128/jb.172.8.4238-4246.1990

Wang YY, Liu B, Zhang XY, Zhou QM, Zhang T, Li H, Yu YF, Zhang XL, Hao XY, Wang M, Wang L, Wei JC. 2014. Genome characteristics reveal the impact of lichenization on lichen-forming fungus *Endocarpon pusillum* Hedwig (*Verrucariales, Ascomycota*). BMC Genomics 15:34 [18 p.]. https://doi.org/10.1186/1471-2164-15-34

Wang YY, Zhang XY, Zhou QM, Zhang XL, Wei JC. 2015. Comparative transcriptome analysis of the lichen-forming fungus *Endocarpon pusillum* elucidates its drought adaptation mechanisms. Science China Life Sciences 58: 89–100. https://doi.org/10.1007/s11427-014-4760-9

White T, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis & al. (eds). PCR protocols: a guide to methods and applications. San Diego, CA. Academic Press. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Yang J, Wei JC. 2008. The new lichen species *Endocarpon crystallinum* from semiarid deserts in China. Mycotaxon 106: 445–448.

Zhang T, Wei JC. 2011. Survival analyses of symbionts isolated from *Endocarpon pusillum* Hedwig to desiccation and starvation stress. Science China Life Sciences 54(5): 480–489. https://doi.org/10.1007/s11427-011-4164-z

Zhang T, Liu M, Wang YY, Wang ZJ, Wei XL, Wei JC. 2017a. Two new species of *Endocarpon* (*Verrucariaceae, Ascomycota*) from China. Scientific Reports 7:7193 [11 p.]. https://doi.org/10.1038/s41598-017-07778-5

Zhang YL, Li H, Wang YY, Wei JC. 2017b. The calcium-binding protein EpANN from the lichenized fungus *Endocarpon pusillum* enhances stress tolerance in yeast and plants, Fungal Genetics and Biology 108: 36–43. https://doi.org/10.1016/j.fgb.2017.09.003